

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ARNT/HIF-1 β in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human ARNT/HIF-1 β Asp517-Ala691 Accession # P27540
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Knockout Validated	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

ARNT (Aryl hydrocarbon nuclear translocator; also HIF-1 β , Hypoxia-inducible factor 1-beta) is an 87-97 kDa member of the bHLH-PAS subclass of the bHLH family of transcription factors. It is widely expressed, and serves as part of both the NF κ B and aryl hydrocarbon receptor (AHR) signaling nodes. For the AHR, the AHR is normally inactive and bound to hsp90 in the cytosol. Following ligand binding, the AHR complex enters the nucleus, dissociates from hsp90, dimerizes with ARNT, and then binds response elements of select genes. Human ARNT is 789 amino acids (aa) in length. It contains a DNA-binding region (aa 87-143), two PAS domains (aa 161-235 and 362-458) and one PAC region (aa 424-467). There are multiple potential splice forms. Two alternate start sites are found at Met10 and Met354, there is a deletion of aa 329-755 and 77-91, and an Ala substitution for aa 328-789.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.