

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Phospho-LKB1/STK11 (S428) in Western blots.
Source	Polyclonal Rabbit IgG
Purification	Antigen Affinity-purified
Immunogen	Phosphopeptide containing human LKB1/STK11 S428 site Accession # Q15831
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

LKB1 (Liver kinase B1; also STK11/Ser/Thr-protein kinase 11 and NY-REN19) is a 55 kDa intracellular member of the LKB1 subfamily, CAMK Ser/Thr protein kinase family of molecules. It is widely expressed, being particularly investigated in small intestinal columnar epithelium and hepatocytes. STK11 is associated with a wide variety of functions, including the initiation of apoptosis through binding to p53, the regulation of TGF-β signaling through the creation of a SMAD4-LIP1-STK11 complex, and the generation of an epithelial polarized phenotype via cytoskeletal remodeling. Schematically, it would appear that STK11 acts, at least in part, through its ability to phosphorylate multiple AMPK-related kinases, as well as AMPK itself. Human STK11 is 433 amino acids (aa) in length. It potentially contains a three aa propeptide at its C-terminus. The mature segment (aa 1-430) possesses one large protein kinase domain (aa 49-309) plus at least four utilized phosphorylation sites. Phosphorylation on Ser428 promotes the ability of LKB1 to suppress G361 cell growth. Palmitoylation is suggested to occur on Cys418. There are three alternative splice variants. Either individually or in combination, they involve an insertion of nine aa after Tyr126 and a 34 aa substitution for aa 371-433. Full-length human STK11 shares 90% aa sequence identity with mouse STK11. Over the area correspondent to the antigen, human and mouse LKB1 are identical in aa sequence.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.