

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Syntaxin 16 in direct ELISAs and Western blots. In direct ELISAs, less than 5% cross-reactivity with recombinant human (rh) Syntaxin 1A, rhSyntaxin 6, and rhSyntaxin BP1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Syntaxin 16 isoform B Leu165-Lys301 Accession # O14662
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Syntaxin 16 (STX16; also SNY16) is a 39 kDa member of the syntaxin family of proteins. It is ubiquitously expressed, embedded in the Golgi membrane, and participates in the fusion of early endosomes with the Golgi stacks. Syntaxin 16 contributes one of four coiled-coil domains necessary for retrograde transport. Human Syntaxin 16 is a type IV single-pass transmembrane protein (very short luminal C-terminus) that is 325 amino acids (aa) in length. It contains a cytoplasmic syntaxin region (aa 74-180), a coiled-coil region (aa 230-292), and a short, three amino acid C-terminal luminal sequence. There are multiple potential isoforms. Two show alternate start sites at Met187 and Met54, while three others show deletions of aa 45-48, 28-48 and 28-44, respectively. A cytoplasmic form shows an Ala substitution for aa 132-325. Over aa 165-301, human Syntaxin 16 shares 95% aa identity with mouse Syntaxin 16.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.