

DESCRIPTION


Species Reactivity	Cotton Rat
Specificity	Detects cotton rat IFN- γ in ELISAs and Western blots. In sandwich immunoassays, less than 0.25% cross-reactivity with recombinant canine IFN- γ , recombinant feline IFN- γ , and recombinant human IFN- γ , recombinant mouse IFN- γ , recombinant rat IFN- γ , and recombinant porcine IFN- γ is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant cotton rat IFN- γ His21-Ile170 Accession # Q9QXX2
Endotoxin Level	<0.10 EU per 1 μ g of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 μ g/mL	Recombinant Cotton Rat IFN- γ (Catalog # 565-RG)
Cotton Rat IFN-γ Sandwich Immunoassay		Reagent
ELISA Capture	0.2-0.8 μ g/mL	Cotton Rat IFN- γ Antibody (Catalog # AF565)
ELISA Detection	0.1-0.4 μ g/mL	Cotton Rat IFN- γ Biotinylated Antibody (Catalog # BAF565)
Standard		Recombinant Cotton Rat IFN- γ (Catalog # 565-RG)
Neutralization		Measured by its ability to neutralize IFN- γ inhibition of EMC-induced cytopathy in the LCRT cotton rat fibrosarcoma cell line. The Neutralization Dose (ND ₅₀) is typically 0.3-1.2 μ g/mL in the presence of 20 ng/mL Recombinant Cotton Rat IFN- γ .

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Interferon-gamma (IFN- γ), also known as type II or immune interferon, exerts a wide range of immunoregulatory activities and is considered to be the prototype proinflammatory cytokine (1, 2). Mature cotton rat IFN- γ exists as a noncovalently linked homodimer of 20-25 kDa variably glycosylated subunits (3). It shares 43%-50% amino acid sequence identity with bovine, canine, equine, feline, human, mouse, rat, rhesus, and porcine IFN- γ . IFN- γ dimers bind to IFN- γ RI (alpha subunits) which then interact with IFN- γ RII (beta subunits) to form the functional receptor complex of two α and two β subunits. Inclusion of IFN- γ RII increases the binding affinity for ligand and the efficiency of signal transduction (4, 5). IFN- γ is produced by a variety of immune cells under inflammatory conditions, notably by T cells and NK cells (6). It plays a key role in host defense by promoting the development and activation of Th1 cells, chemoattraction and activation of monocytes and macrophages, upregulation of antigen presentation molecules, and immunoglobulin class switching in B cells. It also exhibits antiviral, antiproliferative, and apoptotic effects (6, 7). In addition, IFN- γ functions as an anti-inflammatory mediator by promoting the development of regulatory T cells and inhibiting Th17 cell differentiation (8, 9). The pleiotropic effects of IFN- γ contribute to the development of multiple aspects of atherosclerosis (7).

References:

1. Billiau, A. and P. Matthys (2009) *Cytokine Growth Factor Rev.* **20**:97.
2. Pestka, S. *et al.* (2004) *Immunol. Rev.* **202**:8.
3. Houard, S. *et al.* (1999) *Gene* **240**:107.
4. Marsters, S.A. *et al.* (1995) *Proc. Natl. Acad. Sci.* **92**:5401.
5. Krause, C.D. *et al.* (2000) *J. Biol. Chem.* **275**:22995.
6. Schroder, K. *et al.* (2004) *J. Leukoc. Biol.* **75**:163.
7. McLaren, J.E. and D.P. Ramji (2009) *Cytokine Growth Factor Rev.* **20**:125.
8. Muhl, H. and J. Pfeilschifter (2003) *Int. Immunopharmacol.* **3**:1247.
9. Kelchtermans, H. *et al.* (2008) *Trends Immunol.* **29**:479.