

DESCRIPTION

Species Reactivity	Mouse/Rat
Specificity	Detects mouse and rat Neuroglycan C/CSPG5 in direct ELISAs and Western blots. In direct ELISAs, approximately 25% cross-reactivity with recombinant human Neuroglycan C is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse Neuroglycan C/CSPG5 Val31-Gln420 Accession # AAH55736
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Neuroglycan C (NGC; also CSPG5 and CALEB) is a 120-150 kDa type I transmembrane glycoprotein and member of the neuregulin family of proteins (1-2). Depending on its expression profile, NGC may be a glycoprotein of 120 kDa, or a chondroitin sulfate (CS) proteoglycan of 150 kDa (2-3). Mouse NGC is synthesized as a 566 amino acid (aa) precursor that contains a 30 aa signal sequence, a 393 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 122 aa cytoplasmic region. The ECD contains one CS attachment domain (aa 32-273), with CS attachment at Ser117, one EGF-like domain (aa 371-413), three potential sites for N-linked glycosylation, and ten potential sites for O-linked glycosylation (4). Splicing variants produce four isoforms for human NGC. Isoform 1 is the standard form. Isoform 2 has a deletion of aa 487-513, while isoform 3 has an alternative start site at Met82 and the same deletion. Isoform 4 has a 56 aa substitution for aa 514-566. Phosphorylation likely occurs at Ser249, and proteolysis generates a 75 kDa soluble fragment (5). Over aa 31-420, mouse NGC shares 84% aa identity with human NGC. NGC is expressed in nervous tissue and is found on retinal ganglion cells, cerebellar Purkinje cells and hippocampal neurons (6). NGC may function as a growth and differentiation factor involved in neuritogenesis. One study shows that the recombinant ectodomain of NGC core protein enhances neurite outgrowth from rat neocortical neurons in culture via phosphatidylinositol 3-kinase and protein kinase C signaling pathways (7). Another study states that NGC is a novel component of midkine receptors, a heparin-binding growth factor that promotes cell attachment and process extension in oligodendroglial precursor-like cells (3). NGC also acts as a growth factor by directly binding ErbB3 tyrosine kinase and transactivating ErbB2 (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.