

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects human and mouse Bub3 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human Bub1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Bub3 Met1-Thr328 Accession # O43684
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Bub3 (Budding uninhibited by benzimidazoles 3 homolog; also Mitotic checkpoint protein Bub3) is a 35-40 kDa member of the WD repeat Bub3 family of molecules. It plays a key role in chromosome segregation by serving as a component of the mitotic checkpoint complex. This complex inhibits Cdc20, and E3 ubiquitin ligase that initiates the metaphase to anaphase transition, until microtubules are properly aligned with, and bound to, chromatid kinetochores. Human Bub3 is 328 amino acids (aa) in length. It contains five WD40 repeats (aa 5-43, 46-83, 86-124, 128-163 and 223-262) plus a phosphorylation site at Ser211 and a Lys at # 216 that may form an interchain crosslink with Gly. There are two splice variants that show two or four aa deletion over the four C-terminal amino acids. Full-length human and mouse Bub3 share 99% aa identity.