

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse Bub3 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human Bub1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Bub3 Met1-Thr328 Accession # O43684
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Bub3 (Budding uninhibited by benzimidazoles 3 homolog; also Mitotic checkpoint protein Bub3) is a 35-40 kDa member of the WD repeat Bub3 family of molecules. It plays a key role in chromosome segregation by serving as a component of the mitotic checkpoint complex. This complex inhibits Cdc20, and E3 ubiquitin ligase that initiates the metaphase to anaphase transition, until microtubules are properly aligned with, and bound to, chromatid kinetochores. Human Bub3 is 328 amino acids (aa) in length. It contains five WD40 repeats (aa 5-43, 46-83, 86-124, 128-163 and 223-262) plus a phosphorylation site at Ser211 and a Lys at # 216 that may form an interchain crosslink with Gly. There are two splice variants that show two or four aa deletion over the four C-terminal amino acids. Full-length human and mouse Bub3 share 99% aa identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.