

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human WT1 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human WT1 Met127-Gly249 Accession # P19544
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Intracellular Staining by Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

WT1 (Wilms' tumor protein 1; also WT33) is a 52-54 kDa, nuclear member of the EGR C2H2-type zinc-finger family of proteins. Although its predicted MW is 49 kDa, it runs anomalously in SDS-PAGE, likely due to a high proline content. It is widely expressed, being found in developing Sertoli cells, glomerular podocytes, neurons, and CD34+ stem cells. Human WT1 is 449 amino acids (aa) in length. It contains a Pro-rich domain (aa 27-83) and four consecutive C2H2 zinc finger regions (aa 323-347; 353-377; 383-405; 414-438). WT1 forms homodimers, and interacts with multiple molecules. Interaction with the zinc fingers generally promotes gene transcription, while N-terminal interactions block gene transcription. There are at least two dozen splice variants. Some are combinations of deletions of aa 250-266 and 408-410, plus an alternate start site 68 aa upstream of the standard site, and a three aa substitution for aa 1-147. Over aa 127-249, human WT1 shares 98% aa identity with mouse WT1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.