

#### DESCRIPTION


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects endogenous mouse BACH1 in Western blots.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant mouse BACH1 Ser133-Gly513 Accession # P97302
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below

#### DATA



#### PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

#### BACKGROUND

BACH1 (BTB and CNC homolog 1; also basic leucine zipper transcription factor 1) is a ubiquitously expressed member of the Bach family of transcription factors. Although its predicted MW is 82 kDa, it runs anomalously at 110 kDa in SDS-PAGE. BACH1 forms noncovalent homodimers, and heterodimers with Maf oncoproteins and p53-related proteins. It apparently serves as an architectural component for gene regulatory protein complexes. Mouse BACH1 is 739 amino acids (aa) in length. It contains a protein-interaction BTB domain (aa 24-127), a DNA-binding motif (aa 565-580), and a Leu-zipper domain (aa 588-610). This molecule should not be confused with Bach1/Fancj/Brip1 helicase. Over aa 133-513, mouse BACH1 shares 75% and 88% aa identity with human and rat BACH1, respectively.