

DESCRIPTION

Species Reactivity	Human
Specificity	Detects endogenous human ZBTB38 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human ZBTB38 Met1-Tyr232 Accession # Q8NAP3
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

ZBTB38 (Zinc finger and BTB domain-containing protein 38; also CIBZ and Zenon) is an early member of a family of Kaiso-like DNA binding proteins. Although its predicted MW is 134 kDa, it runs anomalously in SDS-PAGE at 175 kDa. ZBTB38 binds to singly methylated CpG (cytosine-phosphate-guanine) motifs. It is considered a transcriptional repressor that either homodimerizes, or heterodimerizes with ZBTB4, and subsequently interacts with either CtBP-1 or -2 to mediate repression. Human ZBTB38 is 1195 amino acids (aa) in length. It contains a BTB domain (aa 33-100), an RD2 region (aa 158-341) and ten C2H2-type zinc finger domains (aa 342-1147). There is one potential alternate start site at Met595. Over aa 1-232, human ZBTB38 shares 96% aa identity with mouse ZBTB38.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.