

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human 5'-Nucleotidase/CD73 in direct ELISAs and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant mouse CD73 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human 5'-Nucleotidase/CD73 Trp27-Lys547 Accession # AAH65937
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

5'-Nucleotidase (also [ecto]-5'-nucleotidase/5'-NT, designated CD73) is a variably glycosylated, 69-73 kDa member of the 5'-Nucleotidase family of enzymes. It is expressed on multiple cell types, including vascular endothelium, transitional and nonkeratinized epithelium, cardiomyocytes, small intestine epithelium, FoxP3⁺ Treg lymphocytes, FDCs and B cells. 5'-Nucleotidase hydrolyzes AMP to adenosine and phosphate. This creates diffusible nucleosides necessary for cell homeostasis, and a ligand for cell membrane adenosine receptors. Mature human 5'-Nucleotidase is a 523 amino acid (aa) GPI-linked protein (aa 27-549). It contains a large Zn-dependent nucleotidase catalytic region (aa 28-532) and a C-terminal substrate binding site (aa 500-506). On the cell surface it exists as a disulfide-linked homodimer. Two splice variants are reported. One shows a deletion of aa 405-454, and a second possesses a 12 aa substitution for aa 253-574. Over aa 1-511, human 5'-Nucleotidase shares 88% aa identity with both mouse and rat 5'-Nucleotidase.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.