

#### DESCRIPTION

<b>Species Reactivity</b>	Mouse/Rat
<b>Specificity</b>	Detects mouse and rat Neurocan in direct ELISAs and Western blots.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant mouse Neurocan Asp23-Asp637 Accession # NP_031815
<b>Conjugate</b>	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
<b>Formulation</b>	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

<b>Western Blot</b>	Optimal dilution of this antibody should be experimentally determined.
<b>Immunohistochemistry</b>	Optimal dilution of this antibody should be experimentally determined.

#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

#### BACKGROUND

Neurocan is a 220 kDa nervous tissue-specific chondroitin sulfate proteoglycan member of the aggrecan/versican proteoglycan family (1). Mouse Neurocan is synthesized as a 1268 amino acid (aa) precursor that contains a 22 aa signal sequence and a 1246 aa mature chain. The mature chain contains one Ig-like V-type domain (aa 37-157), two Link domains (aa 159-254 and 258-356), two EGF-like domains (aa 960-996 and 998-1034), one C-type lectin-like domain (aa 1036-1165), one Sushi domain (aa 1165-1224), and five potential sites for N-linked glycosylation. Mature mouse Neurocan is 90% and 66% aa identical to mature rat and human Neurocan, respectively. Neurocan binds with high affinity to the cell adhesion molecules (CAM) Ng-CAM and N-CAM to inhibit Neuronal adhesion and neurite growth (2-3). In the developing rat retina, the expression of Neurocan is regulated both temporally and spatially, which suggests that it may play a role in the differentiation of and neural network formation of the mammalian retina (1). Injury to the CNS leads to permanent loss of function due to the inability of severed nerve fibers to regenerate back to their targets (4). The lack of CNS repair is attributed in part to the extracellular matrix chondroitin sulfate proteoglycans, such as Neurocan, which are produced by activated glial cells post-injury (4).

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.