

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Pro-Corticoliberin/Pro-CRH in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Pro-Corticoliberin/Pro-CRH Leu25-Lys196 Accession # P06850
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Pro-Corticoliberin (also Pro-CRF and Pro-CRH) is a predicted 19 kDa member of the CRF/Urotensin I family of hormones. It is expressed by placental syncytiotrophoblasts, zona fasciculata adrenal cells and paraventricular neurons, and is induced by glucocorticoids. Pro-Corticoliberin is the precursor to a mature, 41 amino acid (aa) amidated Corticoliberin that induces pituitary ACTH and subsequent cortisol production by the adrenal. Human Pro-Corticoliberin is 170 aa in length (aa 25-194). If unprocessed, it undergoes N-terminal amidation to generate a polypeptide 168 aa in length (aa 27-194). Secreted Pro-Corticoliberin is both bioactive and binds to 37 kDa Corticoliberin-PB. In the placenta, Pro-Corticoliberin is only occasionally cleaved to generate mature Corticoliberin (aa 152-192); in the brain, it undergoes active internal proteolysis to generate multiple fragments, including the 5 kDa mature peptide. Over aa 25-196, human Pro-Corticoliberin shares 78% aa identity with mouse Pro-Corticoliberin.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.