

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse COX-4I1 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human COX4-I1 Ala23-Lys169 Accession # P13073
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cytochrome c oxidase subunit 4 isoform 1 (COX-4I1) is a 21-22 kDa member of the cytochrome c oxidase IV family of proteins. It is a component of COX, an inner mitochondrial membrane multimeric dimer that catalyzes the transfer of electrons from cytochrome c to dioxygen. COX-4I1 is the largest of 10 distinct nuclear DNA-encoded subunits that form each COX monomer. Human COX4-I1 is 169 amino acids (aa) in length and possesses a mitochondrial transit peptide between aa 1-22, an ATP binding site (aa 42; 95-100), and multiple subunit interface sequences. The ATP binding site is suggested to make COX-4I1 a regulatory subunit within the COX complex. There are two potential splice variants for COX-4I1 that show a three aa substitution for aa 81-169, and a single Gly substitution for aa 125-169. COX-4I2, the product of a different gene, shares only 50% aa identity with COX-4I1. Over amino acid 23-169, human COX4-1 shares 78% aa identity with mouse COX-4I1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.