

DESCRIPTION

Species Reactivity	Canine
Specificity	Detects canine IL-21 in direct ELISAs. In direct ELISAs, approximately 35% cross-reactivity with recombinant human IL-21 is observed, and less than 3% cross-reactivity with recombinant mouse IL-21 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant canine IL-21 His18-Ser146 Accession # NP_001003347
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin-21 (IL-21) is a 14 kDa (predicted) four-helix-bundle cytokine and member of the IL-15/IL-21 family. It is made by activated CD4⁺ T cells, activated NKT cells, T helper (Th) cells, and Th17 cells (1-5). Canine IL-21 is synthesized as a 146 amino acid (aa) precursor that contains a 17 aa signal sequence and a 129 aa mature chain. Mature canine IL-21 is 86%, 73%, 66%, and 64% aa identical to mature bovine, human, mouse and rat IL-21, respectively. IL-21 binds to a heterodimeric receptor, which is formed by the common γ -chain subunit (CD134), shared with IL-2, IL-4, IL-7, IL-9, IL-13, and IL-15 receptors, and its own receptor, IL-21 R, a member of the class I cytokine receptor family (1-2). IL-21 R is expressed on a variety of immune cells, including B cells, T cells, dendritic cells, and NK cells, as well as non-immune cells, such as fibroblasts, epithelial cells, and endothelial cells (1-2, 5). Binding of IL-21 to its receptor leads to the activation of members of the JAK-family protein tyrosine kinases, JAK1 and JAK3, and signal transducer and activator of transcription (Stat) molecules (1). IL-21 regulates activation, proliferation, and survival of both CD4⁺ T cells and B cells, the functional activity of CD8⁺ T cells and NK cells, and limits the differentiation of inducible regulatory T cells and counteracts their suppressive properties on effector T cells (1, 5-6). IL-21 also negatively regulates the maturation and function of dendritic cells (1, 5-6). IL-21 also plays a role in many inflammatory and autoimmune diseases, such as Crohn's disease, *Helicobacter pylori*-related gastritis, celiac disease, type I diabetes mellitus, psoriasis, rheumatoid arthritis, and systemic lupus erythematosus (1). IL-21 is also involved in controlling chronic viral infections (7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.