

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human Androgen R/NR3C4 in direct ELISAs. Detects human, mouse and rat Androgen R/NR3C4 in Western blotting.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Androgen R/NR3C4 Thr660-Gln919 Accession # P10275
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The ligand binding domain of human AR (aa 661-920) shares 100% aa sequence identity with mouse and rat AR. AR (Androgen receptor) is a 99 kDa (predicted) member of the NR3 subfamily, nuclear hormone receptor family of proteins. Due to a high number of Gln and Pro residues, it runs anomalously at 100-120 kDa in SDS-PAGE. It is widely expressed, being found in neurons, endothelial cells, osteoblasts, chondrocytes, macrophages, adipocytes, and prostate epithelium. Human AR is 919 amino acids (aa) in length. It contains three discrete domains: a "modulating" N-terminus (aa 1-553) that is rich in Gln, Pro and Gly, a Zn-finger DNA-binding region (aa 554-635), and a ligand-binding domain (aa 672-917). AR is highly polymorphic at the N-terminus, with total Gln and Gly residues differing by seven or more residues among individuals. Multiple potential splice forms exist, including an alternative start site at Met189 and a seven aa substitution for aa 1-538 that generates a 45 kDa isoform. AR does homodimerize, apparently with multiple isotypes. Over aa 661-920, human and mouse are identical in aa sequence.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.