

# **Human LNK/SH2B3 Antibody**

Antigen Affinity-purified Polyclonal Sheep IgG Catalog Number: AF5888


DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human LNK/SH2B3 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	E. coli-derived recombinant human LNK/SH2B3 Gln427-Leu575 Accession # Q9UQQ2
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

#### APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 μg/mL	See Below

#### DATA



Detection of Human LNK/SH2B3 by Western Blot. Western blot shows lysates of HepG2 human hepatocellular carcinoma cell line, SH-SY5Y human neuroblastoma cell line, Ramos human Burkitt's lymphoma cell line, MOLT-4 human acute lymphoblastic leukemia cell line, and A431 human epithelial carcinoma cell line. PVDF membrane was probed with 0.5 µg/mL of Sheep Anti-Human LNK/SH2B3 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF5888) followed by HRPconjugated Anti-Sheep IgG Secondary Antibody (Catalog # HAF016). A specific band was detected for LNK/SH2B3 at approximately 75 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.

PREPARATION AND STORAGE		
Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.	
Shipping	pping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.	
	*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C	

## Stability & Storage

19 -

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 6 months, -20 to -70 °C under sterile conditions after reconstitution.

### BACKGROUND

LNK (also SH2B adaptor protein 3) is a protein intermediary, or LiNK, between cell surface molecules (TPO R; EPO R; SCFR; TCR) and downstream signaling molecules such as Pl3 K, Grb2 and Jak2. Although its predicted MW is 63 kDa, it runs anomalously at 75 kDa in SDS-PAGE. LNK is found in hematopoietic cells, and serves to limit responsiveness to growth factor stimulation. Human LNK is 575 amino acids (aa) in length. It contains an N-terminal Phe-zipper segment that likely serves as a dimerization domain (aa 24-81), a PH domain that may interact with membrane lipids (aa 197-305), and an SH2 domain that binds phosphoTyr (aa 364-462).

Rev. 2/6/2018 Page 1 of 1

