

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Contactin-6 in direct ELISAs and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant human (rh) Contactin-6 is observed, and less than 5% cross-reactivity with recombinant mouse (rm) Contactin-3, rmConta
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse Contactin-6 Glu20-Ser998 Accession # Q9JMB8
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Contactin-6, also known as NB-3, is a 130 kDa member of the TAG/F3 subgroup of Contactin GPI-linked neuronal adhesion proteins (1, 2). Mature mouse Contactin-6 consists of six immunoglobulin-like domains, a flexible linker region, four fibronectin type III domains, and a GPI anchor (3-5). Mouse Contactin-6 shares 90% and 96% amino acid (aa) sequence identity with human and rat Contactin-6, respectively. Alternate splicing generates an isoform with a 17 aa deletion in the first Ig-like domain (3). Contactin-6 expression is upregulated postnatally in the cerebellum, brainstem, and hippocampus, and it shows differential expression within various structures of the brain (3, 5-7). In the cerebellum, it is presynaptically localized at synapses between excitatory glutamatergic parallel fibers and Purkinje cells (7). Contactin-6 associates *in cis* with CHL-1 and *in trans* with Notch-1 on oligodendrocytes (6, 8). Its binding to Notch-1 triggers translocation of the Notch intracellular domain to the nucleus and promotes oligodendrogenesis (6). Contactin-6 function is important in the postnatal development and function of the cerebellum, as shown in knockout mice which exhibit poor motor coordination (5, 7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.