

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Lipase I in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Lipase I Arg14-His154 Accession # Q6XZB0
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Lipase I (Lipase member I; also mPA-PLA₁βG LPDL and PLA1C) is a 55 kDa member of the PLA₁ subfamily, pancreatic lipase family, AB hydrolase superfamily of enzymes. It is expressed by sperm, where it is apparently secreted and subsequently binds to membrane-associated HSPGs. Lipase I hydrolyzes phosphatidic acid to generate 2-acyl-LPA, which then activates one of four G-protein coupled receptors. Human Lipase I precursor is 460 amino acids (aa) in length. It contains a 15 aa signal sequence, followed by an active catalytic site composed of Ser159, Asp183 and His268. Lipase I is known to be phosphorylated on Ser25. There are two isoform variants, both of which show a 34 aa substitution for aa 1-13, and only one of which shows an eight aa substitution for aa 245-460. Over aa 14-154, human Lipase I shares 61% aa identity with mouse Lipase I.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.