

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CD1c in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) CD1a, rhCD1b, rhCD1d, and rhCD1e is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CD1c/BDCA-1 Ala19-Met302 (Phe300Ser) Accession # P29017
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CD1c (Cluster of differentiation antigen 1c) is a 316 amino acid glycoprotein with a predicted molecular weight of 43 kDa. CD1c is a member of the CD1 family of molecules. It is expressed by thymocytes, dendritic cells and B cells, and exists as part of a noncovalent complex with 12 kDa β_2 -microglobulin. It is found in the plasma membrane and early endosomes (but not lysosomes), and is presumed to present glycolipids and acylated peptides to T cells. Mature human CD1c is a 316 amino acid (aa) type I transmembrane glycoprotein. It contains a 285 aa extracellular domain (ECD) (aa 18-302) plus a 10 aa cytoplasmic tail. The ECD shows one Ig-like domain (aa 203-296) that associates with β_2 -microglobulin, and a TyrGlnAspIle internalization motif in the cytoplasmic tail. There are three potential splice variants. One shows a Trp substitution for aa 327-333, a second shows an eight aa substitution for aa 298-333, and a third shows a 50 aa substitution for aa 297-333. There appears to be no direct mouse counterpart to human CD1c.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.