

DESCRIPTION

Species Reactivity	Mouse/Rat
Specificity	Detects mouse and rat BMP-15/GDF-9B in Western blots and recombinant mouse in direct ELISAs. In direct ELISAs, approximately 5% cross-reactivity with recombinant human (rh) BMP-15 is observed and less than 1% cross-reactivity with rhBMP-3 a
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse BMP-15/GDF-9B Gln268-Arg392 Accession # Q9Z0L4
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

BMP-15 (Bone morphogenetic protein 15; also GDF-9B) is a 24-26 kDa member of the TGF- β superfamily of proteins. It is expressed by late primary follicle oocytes, where it promotes the transition of preantral granulosa cells to cumulus cells, and later the expansion of cumulus cells. Mouse BMP-15 precursor is a 50-55 kDa, 367 amino acid (aa) glycoprotein. It is proteolytically cleaved to generate a 40 kDa prosegment (aa 26-267) plus a 24 kDa, 124 aa mature region (aa 268-392) that may be phosphorylated (on Ser7 of the mature molecule) and/or glycosylated. Secreted BMP-15 does not occur as a mature homodimer, but it does exist as a mature monomer, an uncleaved precursor, or as a noncovalent heterodimer composed of a cleaved mature region and its prosegment. The heterodimer may also form an oligomer. The BMP-15 prosegment reportedly forms a noncovalent heterodimer with 20 kDa mature GDF-9. Mature mouse BMP-15 shares 70% and 91% aa identity with human and rat BMP-15, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.