

DESCRIPTION

Species Reactivity	Mouse/Rat
Specificity	Detects recombinant mouse Carboxylesterase 3/CES3 in direct ELISAs and Western blots. Detects intrinsic CES3 in mouse and rat tissues in Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) CES1, rhCES2, recombinant m
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Carboxylesterase 3/CES3 Tyr19-Glu561 Accession # Q8VCT4
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Carboxylesterase 3 is a member of a large family of carboxylesterases that are responsible for the hydrolysis of ester and amide bonds (1, 2). These carboxylesterases are widely distributed in mammalian tissues and have broad substrate specificity ranging from small molecule esters such as phenylester to long-chain fatty acid esters and thioesters. Because many ester-containing drugs require the ester linkage to improve the bioavailability of therapeutic agents, CESs play a major role in drug metabolism and activation. CES3 is also known as triacylglycerol hydrolase for its lipolytic activity. It has been reported to be the major lipase in white adipose tissue (3). Mouse CES3 shares 93% homology with rat CES3.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.