

DESCRIPTION

Species Reactivity	Mouse/Rat
Specificity	Detects mouse and rat CCR8 in direct ELISAs and Western blots. In direct ELISAs, less than 2% cross-reactivity with recombinant mouse (rm) CCR1, rmCCR4, and rmCCR5 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse CCR8 Met1-Met33, Asn92-Lys105, Tyr170-Ile200, His262-Leu278 Accession # P56484
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CCR8 (C-C chemokine receptor 8; also CD198) is a 41-43 kDa member of the GPCR #1 family of transmembrane proteins. Mouse CCR8 is expressed on vascular smooth muscle cells, monocytes, eosinophils, peritoneal macrophages, thymocytes, CD8⁺ T cells, Langerhans cells and neurons. CCL1/TCA3 and vMIP-1 are known agonists for CCR8. Mouse CCR8 is a 7-transmembrane protein that is 353 amino acids (aa) in length. It contains a 33 aa N-terminal extracellular domain plus a 50 aa C-terminal cytoplasmic tail. In mouse, CCR8 is N- and possibly O-glycosylated, and known to be sulfated on Tyr14 and 15. The unusual nature of these posttranslational modifications may lead to anomalous migration in SDS-PAGE. Over aa sequences 1-33 and 92-105 collectively, mouse CCR8 shares 64% and 85% aa identity with human and rat CCR8, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.