

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat Jagged 1 in ELISAs and Western blots. In sandwich ELISAs, approximately 15% cross-reactivity with recombinant human (rh) Jagged 1 is observed, and less than 0.3% cross-reactivity with rhJagged 2 and recombinant mouse Jagged 2 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant rat Jagged 1 Ser32-Asp1068 (Gly57-Arg59 del, Asp63Thr, Arg64Leu, and Val-Arg-Pro-Tyr ins before Lys69) Accession # Q63722
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Rat Jagged 1 was the first ligand for Notch identified in mammals. Since both the ligands and receptors are transmembrane proteins, direct cell-cell interactions are thought to be required for activating Notch signaling. Jagged 1 is synthesized as a precursor protein that contains a 21 aa signal sequence, a 1048 aa extracellular region, a 25 aa transmembrane (TM) segment and a short, 226 aa cytoplasmic domain. The large extracellular region has a DSL (Delta, Serrate, Lag-2 consensus sequence) domain followed by 16 EGF-like repeats, and a cysteine-rich (CR) region (1). The extracellular region of rJagged 1 binds to multiple Notch receptors on the cell surface as well as in solid phase binding studies. The DSL motif is necessary for binding to Notch receptors and the EGF repeats modulate the affinity of the interaction with Notch receptors (2). Notch signaling is implicated in many developmental processes in a variety of cell types. Jagged-Notch signaling specifies cell fate, regulates pattern formation, defines boundaries between different cell types, and modulates cell proliferation and differentiation. Some specific areas where Jagged is involved include hematopoiesis, myogenesis, neurogenesis and development of the vasculature (3). For instance soluble, non-transmembrane forms of Jagged 1 influence behavior in fibroblast cells leading to characteristics exhibited by endothelial cells during angiogenesis (4). Soluble Jagged 1 is also capable of maintaining the survival and enhancing the expansion of human stem cells that are capable of reconstituting hematopoietic lineages *in vivo* (5). Furthermore, Jagged 1 is implicated in human disease: Alagille syndrome, an autosomal dominant disorder characterized by defects in liver, heart, eye, skeletal, craniofacial tissues, and kidney, is caused by mutations in Jagged 1 (6). Depending on cell types and how soluble forms of the ligand are presented, ligand binding can result in activation or inhibition of Notch signaling (7). Rat Jagged 1 shows 98% and 99% aa identity to human and mouse Jagged 1 extracellular domains respectively. Relative to the extracellular region of rat Jagged 2, the aa identity is 58%.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.