

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse Cyclin B2 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse Cyclin B2 Ala2-Leu101 Accession # P30276
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cyclin B2 (also CCNB2 and G2/mitotic-specific Cyclin B2) is a 48-54 kDa member of the cyclin AB subfamily, cyclin family of proteins. It is widely expressed and associates with CDK1. Here, it provides substrate specificity to a phosphorylating complex. A phosphor-CDK1:Cyclin B2 complex is associated with the Golgi apparatus, where it contributes to Golgi fragmentation during mitosis. It is also possible that Cyclin B2 can substitute for Cyclin B1 during the early stages of mitosis. Human Cyclin B2 is 398 amino acids (aa) in length. It contains two cyclin box folds (aa 201-290 and 298-383) and two substrate binding sites (aa 165-254 and 264-347). Phosphorylation occurs at Ser92, Thr94, Ser99, Ser204, Ser392 and Ser398. Over aa 1-101, human Cyclin B2 shares 79% aa identity with mouse Cyclin B2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.