

DESCRIPTION

Species Reactivity	Viral
Specificity	Detects Viral MIP-II in direct ELISAs and Western blots. In direct ELISAs, less than 10% cross-reactivity with recombinant viral MIP-I and recombinant human MIP-1β is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Viral MIP-II Leu24-Arg94 Accession # NP_572064
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human herpesvirus-8 (HHV-8)/Kaposi's sarcoma-associated herpesvirus (KSHV) is a γ herpesvirus with homology to herpesvirus Saimiri and Epstein-Barr virus. HHV-8 is etiologically linked to Kaposi's sarcoma and a B-cell lymphoma known as primary effusion lymphoma. HHV-8 has been shown to encode a variety of immunomodulatory proteins which were apparently pirated from cellular genes by the virus. Three chemokine-like proteins, vMIP-I, vMIP-II and vMIP-III have been found to be encoded within the HHV-8 genome.

Viral MIP-II cDNA encodes a 94 amino acid (aa) residue precursor protein with a 23 aa residue signal peptide that is cleaved to yield a 71 aa residue mature protein. Among human chemokines, vMIP-II is most closely related to MIP-1α, sharing approximately 41% amino acid sequence identity. At the amino acid sequence level, vMIP-I and vMIP-II also share 48% identity. vMIP-I and vMIP-II are more closely related to one another phylogenetically than to other human chemokines, suggesting that they may have arisen by gene duplication within the virus rather than by two independent gene acquisitions. vMIP-II binds to the CCR-3 chemokine receptor through which eotaxin and other β chemokines activate eosinophils. vMIP-II has been shown to activate and chemoattract human eosinophils. Both vMIP-I and vMIP-II have been shown to partially block HIV infection of peripheral blood mononuclear cells. vMIP-I and vMIP-II have also been found to be highly angiogenic in the chorioallantoic assay, suggesting that they may be partially responsible for the marked vascularity seen in KSHV-associated tumors.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.