

DESCRIPTION

Species Reactivity	Human
Specificity	Detects Carboxypeptidase B2/CPB2 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) CPA1, rhCPB1 and recombinant mouse CPB1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Carboxypeptidase B2/CPB2 Phe23-Val423 (Ala169Thr) Accession # Q96IY4
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CPB2 (Carboxypeptidase B2; also CPU and TAFI) is a secreted, 50-60 kDa glycoprotein member of the peptidase M14 family of enzymes. It is expressed by hepatocytes (60 kDa) and platelets (50 kDa), with MW differences attributable to glycosylation. CPB2 is cleaved by thrombin and plasmin, generating a 36 kDa, relatively insoluble nonglycosylated enzymatically active fragment (TAFIa). Active CPB2 removes C-terminal Lys residues from fibrin, thereby interrupting plasmin generation and promoting fibrin polymerization. Human CPB2 (proprecursor/zymogen) is 401 amino acids (aa) in length. It contains a prosequence (aa 23-114) and an active fragment (aa 115-423) that acts on C-terminal Lys or Arg residues. There is one potential isoform variant that shows a deletion of aa 198-234 accompanied by a 16 aa substitution for aa 382-423. Over aa 23-423, human CPB2 shares 85% aa identity with mouse CPB2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.