

C. botulinum BoNT-D Light Chain Alexa Fluor® 750-conjugated Antibody

Antigen Affinity-purified Polyclonal Sheep IgG Catalog Number: AF6037S 100 µg

DESCRIPTION				
Species Reactivity	C. botulinum			
Specificity	Detects <i>C. botulinum</i> BoNT-D Light Chain in direct ELISAs and Western blots. In direct ELISAs, approximately 5% cross-reactivity with recombinant Light Chains of BoNT-A, -B, -C1, -E, -F, and -G is observed.			
Source	Polyclonal Sheep IgG			
Purification	tion Antigen Affinity-purified			
Immunogen	E. coli-derived recombinant Clostridium BoNT-D Light Chain Thr2-Ser428 Accession # P19321			
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm			
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide			
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.			

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

DE	EDA	DAT	ION.	AND	STO	RAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.		
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied		

BACKGROUND

BoNT-D LC (Botulinum neurotoxin serotype D light chain) is a 50-55 kDa member of the peptidase M27 family of molecules. It is the product of Clostridium botulinum, and inhibits acetylcholine release from neuromuscular junctions. This is accomplished by toxin internalization with subsequent cleavage of synaptobrevin, thus blocking synaptic vesicle fusion with the presynaptic membrane. Notably, BoNT-D is not toxic to human cells. BoNT/D precursor is 1276 amino acids (aa) in length. Following internalization and precursor proteolytic cleavage, it assumes a mature form that shows a disulfide-linked 442 aa light chain/enzyme N-terminus, and a 834 aa heavy chain/receptor-binding C-terminus. The 100 kDa heavy chain creates a channel within the endosome that allows for redox rupture of the disulfide bond and entry of the light chain into the cytosol. Covalent attachment of cargo molecules to the N-terminus is used to transport molecules into the cytosol. The D light chain shares less that 50% aa identity with the other botulinum serotype light chains.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/16/2025 Page 1 of 1

Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956