

Mouse MCAM/CD146 Alexa Fluor® 700-conjugated Antibody

Antigen Affinity-purified Polyclonal Sheep IgG Catalog Number: AF6106N 100 µg

DESCRIPTION		
Species Reactivity	Mouse	
Specificity	Detects mouse MCAM/CD146 in direct ELISAs and Western blots. In direct ELISAs, approximately 40% cross-reactivity with recombinant rat MCAM is observed and approximately 10% cross-reactivity with recombinant human MCAM is observed.	
Source	Polyclonal Sheep IgG	
Purification	Antigen Affinity-purified	
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse MCAM/CD146 Val24-Tyr563 Accession # Q8R2Y2	
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm	
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide	
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.	

APPLICATIONS			
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.			
Western Blot	Optimal dilution of this antibody should be experimentally determined.		
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.		

PREPARATION AND STORAGE		
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied	

BACKGROUND

MCAM (Melanoma cell adhesion molecule; also CD146 and MUC18) is a 110-120 kDa member of a small group of Ig-superfamily molecules that includes CD239 and CD166. MCAM has also been reported at a molecular weight of approximately 150 kDa. In rodent, MCAM is reportedly expressed on neurons, endothelial cells, NK cells, neutrophils, mesenchymal stem cells and melanoma cells. MCAM appears to contribute to intercellular endothelial cell junctions, and possibly contributes to the migration of select cell types. Mature mouse MCAM is a 625 amino acid (aa) type I transmembrane glycoprotein. Its extracellular region is 540 aa in length (aa 24-563). It contains two V-type Ig-like domains (aa 24-244) followed by three C2-type Ig-like domains (aa 246-512). One cytoplasmic region splice form shows a seven aa substitution for aa 600-648. Unlike human, rodent MCAM does not undergo a splicing event that will generate a soluble isoform. Over aa 24-563, mouse MCAM shares 90% and 74% aa identity with rat and human MCAM, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/16/2025 Page 1 of 1

Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956

Bio-Techne®

USA | TEL: 800.343.7475 Canada | TEL: 855.668.8722 Europe | Middle East | Africa TEL: +44.0.1235.529449

China | info.cn@bio-techne.com TEL: 400.821.3475