

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Adrenomedullin/ADM in direct ELISAs.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Adrenomedullin/ADM Ala22-Gly147 Accession # P35318
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Adrenomedullin (ADM; also AM) is a secreted, monomeric, 6 kDa member of the Adrenomedullin family of molecules. It is widely expressed, being found in smooth muscle cells, endothelium, adrenal medulla chromaffin cells, fibroblasts and keratinocytes. ADM has multiple functions, including vasodilation, the maintenance of vascular integrity, and the suppression of inflammatory mediator secretion. The human ADM preproprecursor is 185 amino acids (aa) in length. It contains a 21 aa signal sequence, a processed 20 aa peptide termed PAMP (aa 22-41), an N-terminal propeptide (aa 45-92), the ADM precursor (amidation is required for maturation) (aa 95-147), and a C-terminal propeptide (aa 148-185). The ADM precursor with a terminal Gly147 circulates naturally with bioactive, mature amidated ADM (aa 95-146). Depending upon the tissue, truncated forms of ADM likely also occur, including variants spanning aa 120-146 and 128-146. Over aa 22-147, the human ADM preproprecursor shares 70% aa identity with mouse ADM preproprecursor.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.