

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CCR5 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant mouse (rm) CCR1, rmCCR4, and rmCCR8 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse CCR5 Met1-Ala32, Ala92-Lys104, Thr169-Met200, Gln263-Gln279 (Ile11Ser, Ile97Val) Accession # P51682
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CCR5 (CC chemokine receptor 5; also CD195 and MIP-1α receptor) is a 41kDa (predicted) member of the G-protein coupled receptor #1 family of proteins. It is expressed on Treg T cells, NK cells, neurons, macrophages and Th1 T cells. CCR5 responds to RANTES, MIP-1α and -1β, and induces cell adhesion and migration. Mouse CCR5 is a 7-transmembrane glycoprotein 354 amino acids (aa) in length and it contains an O-glycosylated and sulfated extracellular N-terminus (aa 1-32), plus a phosphorylated and palmitoylated intracellular C-terminus (aa 304-354). In human and primates, CCR5 appears as a 62-64 kDa membrane form, possibly due to extensive posttranslational modification. CCR5 will form homodimers, heterodimers with CCR2, and heterooligomers with CCR2 and CXCR4. Over aa 1-32, 92-104, 169-200 and 263-279 collectively, mouse CCR5 shares 88% and 78% aa identity with rat and human CCR5, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.