

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human BMP-5 in direct ELISAs and Western blots. In direct ELISAs, less than 5% cross-reactivity with recombinant human (rh) BMP-2, rhBMP-4, rhBMP-6, rhBMP-7, and rhBMP-8 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human BMP-5
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Bone Morphogenetic Protein-5 (BMP-5) is one of at least 15 structurally and functionally related BMPs which are members of the transforming growth factor β (TGF- β) superfamily (1). BMP-5 is synthesized as a 454 amino acid (aa) precursor protein that is cleaved at the dibasic cleavage site (RxxR) to release the 20 kDa C-terminal mature protein (2). Mature BMP-5 contains seven conserved cysteine residues involved in formation of the cysteine knot and the single interchain disulfide bond. Biologically active BMP-5 is a disulfide-linked homodimer of the C-terminal mature protein. Mature human BMP-5 shares 96% aa sequence identity with mouse and rat BMP-5. Cellular responses to BMP-5 are mediated by the formation of hetero-oligomeric complexes of type I and type II serine/threonine kinase receptors (1). BMP-5 is expressed by chondrocytes in proliferating and hypertrophic zones of bone growth plates (3). It contributes to limb development by promoting proliferation and differentiation of chondrocytes as well as apoptosis of undifferentiated mesoderm (3, 4). Genetic defects in BMP-5 which cause C-terminal truncation or loss of the proteolytic cleavage site result in multiple skeletal abnormalities, including the *short ear* phenotype in mice (5, 6). BMP-5 is also expressed by ovarian granulosa cells where it functions as an autocrine factor to promote GC proliferation and inhibit their production of progesterone (7). In the nervous system, BMP-5 promotes dendrite outgrowth and dopaminergic neuronal differentiation (8, 9). It is upregulated in oral squamous carcinoma cells and induces the apoptosis of some myeloma cell lines (10, 11).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.