

**DESCRIPTION**



<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human PRDM14 in direct ELISAs and Western blots.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human PRDM14 Ala2-Asp201 Accession # Q9GZV8
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.1 µg/mL	See Below
<b>Immunocytochemistry</b>	5-15 µg/mL	See Below

**DATA**

<p><b>Western Blot</b></p>  <p><b>Detection of Human PRDM14 by Western Blot.</b> Western blot shows lysates of BG01V human embryonic stem cells (undifferentiated) and NTera-2 human testicular embryonic carcinoma cell line. PVDF Membrane was probed with 0.1 µg/mL of Sheep Anti-Human PRDM14 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF6175) followed by HRP-conjugated Anti-Sheep IgG Secondary Antibody (Catalog # HAF016). A specific band was detected for PRDM14 at approximately 65 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p><b>Immunocytochemistry</b></p>  <p><b>PRDM14 in BG01V Human Embryonic Stem Cells.</b> PRDM14 was detected in immersion fixed BG01V human embryonic stem cells using Sheep Anti-Human PRDM14 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF6175) at 10 µg/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 493-conjugated Anti-Sheep IgG Secondary Antibody (green, upper panel; Catalog # NL012) and counterstained with DAPI (blue, lower panel). Specific staining was localized to nuclei. View our protocol for <a href="#">Fluorescent ICC Staining of Stem Cells on Coverslips</a>.</p>
--	---

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

PRDM14 is an approximately 65 kDa transcriptional inhibitor that contains one histone methylating PR/SET domain (aa 266-367) and six zinc finger repeats (aa 400-568). PRDM14 is preferentially expressed in germ cells and undifferentiated embryonic stem cells in which it plays a central role in lineage specification, maintenance of pluripotency, and repression of somatic gene transcription. PRDM14 is up-regulated in breast cancer and lymphoblastic leukemia. Within aa 1-201, human and mouse PRDM14 share 46% aa sequence identity.