

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CD96 in direct ELISAs and Western blots. In direct ELISAs, less than 3% cross-reactivity with recombinant mouse CD96 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CD96 Lys25-Asp501 Accession # NP_005807
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CD96 (also Tactile) is a 160 kDa member of the Ig-Superfamily. It is expressed on CD4⁺ and CD8⁺ T cells, plus NK cells and select B cells. Human CD96 binds to CD155 and presumably participates in NK cell killing of CD155-expressing target cells. Mature human CD96 is a 564 amino acid (aa), type I transmembrane glycoprotein. It contains a 498 aa extracellular region (aa 22-519) that contains three Ig-like domains. The two N-terminal domains are V-type (aa 38-238), while the distal domain is a C-type structure (aa 269-375). There is one isoform that shows a deletion of aa 183-192. This deletion converts the second V-type domain into an I-like domain, and generates the most common form of CD96. An additional isoform shows the same deletion coupled to a nine aa substitution for aa 410-585. Over aa 1-536, human CD96 shares 59% aa identity with mouse CD96.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.