

DESCRIPTION

Species Reactivity	Viral
Specificity	Detects viral CMV UL146/vCXC1 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant viral CMV UL146/vCXC1
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cytomegalovirus (CMV), a member of the beta herpesvirus subfamily, typically causes subclinical or latent infections in the normal adult population. However, CMV can cause congenital disease during pregnancy and is a human opportunistic pathogen that affects immunocompromised individuals. The CMV genome has been shown to contain homologs of cellular immunomodulatory proteins, including US28 (a CC chemokine receptor) and a MHC class I homolog. Virulent CMV clinical isolates have also been shown to carry at least 19 genes, designated *UL133-UL151*, that are not found in laboratory strains that have lost virulence characteristics. Two of these genes, *UL146* and *UL147*, exhibit sequence similarity to CXC chemokines.

The CMV *UL146* open-reading frame encodes a 117 amino acid residue precursor protein with a predicted 22 residues signal peptide that is cleaved to generate the mature protein. Recombinant UL146 has been shown to induce calcium mobilization, chemotaxis and degranulation of neutrophils.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.