

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TRIF/TICAM1 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human TRIF/TICAM1 Lys29-Ala204 Accession # Q8IUC6
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

TRIF (TIR domain-containing adaptor inducing IFN-β; also TICAM1) is a 105-110 kDa cytoplasmic adaptor molecule that mediates Toll receptor signaling. It is widely expressed and associates with both TLR3 and TLR4. Relative to TLR3, TRIF appears to activate IRF 3, -4, and -7, as well as NFκB and FADD. Its action on FADD is through RIP1, and this induces apoptosis. Human TRIF is 712 amino acids (aa) in length and contains three TRAF6 bonding motifs (aa 84-91, 248-255 and 299-309), one TIR domain (aa 390-460), a Pro-rich region (aa 614-678), and an overlapping RHIM domain (aa 661-699). The molecule is reported to form a homodimer. There are multiple potential isoform variants. One shows a 23 aa substitution for aa 31-162 accompanied by a Pro substitution for aa 633-660, a second shows a 44 aa substitution for aa 352-712, and a third shows a 38 aa substitution for aa 271-712. Over aa 29-204, human TRIF shares 62% aa identity with mouse TRIF.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.