

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse and human FGF-10 in Western blots. In direct ELISAs, approximately 100% cross-reactivity with recombinant rat FGF-10 is observed and less than 10% cross-reactivity with recombinant mouse FGF-6 and recombinant mouse F
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse FGF-10 Ser62-Thr209 Accession # NP_032028
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The Fibroblast Growth Factors (FGFs) are heparin binding glycoproteins that exert a variety of biological activities toward cells of mesenchymal, neuronal, and epithelial origin. FGF-10 belongs to the subgroup of FGFs that also includes FGF-3, -7, and -22 (1). Mature mouse FGF-10 is an approximately 20 kDa protein that contains a serine-rich region near its N-terminus (2, 3). C-terminal to this region (aa 62-209), it shares 94% and 100% amino acid sequence identity with human and rat FGF-10, respectively. FGF-10 is secreted by mesenchymal cells and associates with extracellular FGF-BP (1, 4). It preferentially binds and activates epithelial cell FGF R2 (IIIb) and interacts more weakly with FGF R1 (IIIb) (5). The mitogenic and chemotactic properties of FGF-10 are critical in many tissues during embryogenesis. This includes limb bud initiation (6), palate development (7), branching morphogenesis and directional outgrowth of lung buds (2, 8), formation of the otic vesicle and cochlea (9), adipogenesis (10), and the development of prostate, mammary, lacrimal, and submandibular salivary glands (11-14). FGF R2 (IIIb) signaling in these responsive tissues is similarly important during embryogenesis (7, 9, 12-14). The expression and function of FGF-10 are negatively regulated by Shh and BMP-4 in the developing lung (2, 8). Overlapping expression patterns and activities with FGF-3, -7, and -8 suggest at least a partial redundancy in FGF-10 biology (7, 9, 13, 14). FGF-10 induced signaling through FGF R2 (IIIb) also contributes to the progression of pancreatic cancer (15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.