

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects human and mouse DCBLD2/ESDN in Western blots and human DCBLD2/ESDN in direct ELISAs.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human DCBLD2/ESDN Gln67-Ala528 Accession # Q96PD2
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	See Below
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

DCBLD2 (discoidin, CUB, and LCCL domain containing 2), also known as ESDN (endothelial and smooth muscle cell derived neuropilin-like) and CLCP1, was identified through its up-regulation in a highly metastatic lung cancer cell line. DCBLD2 has structural similarities to neuropilins, VEGF receptors, and semaphorins, and is thought to play a role in cell motility and metastasis. SEMA4B is a receptor for DCBLD2/ESDN/CLCP1 and this interaction may be involved in regulating cancer cell metastasis.