

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human TDRD1 in direct ELISAs and Western blots. In direct ELISAs, approximately 25% cross-reactivity with recombinant mouse TDRD1 is observed and less than 1% cross-reactivity with recombinant human TDRKH is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human TDRD1 Asn837-Lys968 Accession # Q9BXT4
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

TDRD1 (Tudor domain containing protein 1; also CT41.1) is a 120-140 kDa member of the Tudor family of proteins. It is expressed in spermatocytes and spermatids, and forms part of an RNA-silencing complex that regulates gene expression. TDRD1, through its Tudor and MYND domains, binds methylated PIWI (P-element-induced wimpy testis) protein and PIWI-interacting RNA loading factor, respectively. This serves to promote the recruitment of germline-specific small RNAs into a PIWI ribonucleoprotein complex. Human TDRD1 is 1180 aa in length. It contains one MYND-type Zn finger region (aa 170-206), and four methyl-binding Tudor domains (aa 312-1048). There are four potential splice variants with predicted molecular weights ranging between 79 kDa and 133 kDa. One contains an alternate start site at Met404, another contains an alternative start site at Met22, accompanied by deletions of aa 554-610 and 775-831, a third shows a 10 aa substitution for Ser1180, while a fourth contains deletions of aa 328-375, 554-610 and 1059-1134. Over aa 837-968, human TDRD1 shares 66% aa identity with mouse TDRD1.