

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human SNAP23 in direct ELISAs and Western blots. In direct ELISAs, approximately 5% cross-reactivity with recombinant mouse SNAP23 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human SNAP23 Asp146-Ser211 Accession # O00161
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SNAP23 (Synaptosomal associated protein 23; also Syndet) is a membrane-associated 23-29 kDa member of the SNAP25 family of proteins. It is expressed in multiple cell types, including neutrophils, platelets, mast cells and adipocytes. SNAP23 is involved in vesicle exocytosis. It has been proposed that SNAP23 can associate with both vesicle and target (cell) membranes via a lipid modification. As a v-SNARE, it may interact with syntaxin-6 at the cell membrane. As a t-SNARE, in conjunction with syntaxin-4, it likely interacts with VAMP-2 and -8 on the vesicle membrane. In either case, this approximates two membranes, which subsequently fuse to create a pore. Human SNAP23 is 211 amino acids (aa) in length. It contains two t-SNARE coiled-coil homology domains (aa 14-76 and 148-207). Palmitoylation occurs between aa 80-87, while phosphorylation occurs on multiple Ser/Thr residues. There is one splice variant that shows a deletion of aa 90-142. Over aa 146-211, human and mouse SNAP23 share 85% aa identity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.