

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat Oncomodulin in direct ELISAs and Western blots. In direct ELISAs, approximately 40% cross-reactivity with recombinant human Oncomodulin is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant rat Oncomodulin Met1-Ser109 Accession # P02631
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Oncomodulin (OM; also parvalbumin beta) is a 12 - 14 kDa member of the parvalbumin family of Ca⁺⁺-binding proteins. It is expressed in early embryonic cells, placenta, and in tumors. OM was originally thought to have expression restricted to neoplastic tissues, early embryonic cells and certain tumor cell lines. Recent research shows that oncomodulin is also expressed and secreted by macrophages where, in the retina, it binds to retinal ganglion cells (RGCs) and functions to promote axon regeneration in early embryonic cells, placenta, and in tumors. OM is both cytoplasmic, and secreted. Rat OM is 109 amino acids (aa) in length. It contains a vestigial Ca⁺⁺-binding site (aa 733) and two EF hand domains, the latter of which contains one high-affinity Ca⁺⁺-binding site (aa 78108). Relative to parvalbumin alpha, OM has a lower pI (<4.8), a higher affinity for Ca⁺⁺, and they share only 50% aa identity. Full length rat OM shares 95% and 89% aa identity with mouse and human OM, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.