

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Collagen XIII in direct ELISAs and Western blots. In direct ELISAs, approximately 40% cross-reactivity with recombinant mouse COL13A1 and less than 1% cross-reactivity with recombinant human (rh) COL23A1 and rhCOL25A1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Collagen XIII Glu109-Leu668 Accession # NP_542992
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

COL13A1 (collagen 13-alpha 1) is a 95 kDa member of the transmembrane group of the collagen family of proteins. It is a type II transmembrane glycoprotein that is expressed by multiple cell types, including fibroblasts, endothelial cells, and cardiac muscle. COL13A1 forms disulfide-linked homotrimers and participates in cell adhesion by binding to the integrin α 1 subunit, nidogen-2, and fibronectin. Human COL13A1 is 717 amino acids (aa) in length (SwissProt #:Q5TAT6). It has an N-terminal 44 amino acid cytoplasmic region plus a 656 amino acid extracellular domain (ECD) (aa 62-717). The ECD contains four non-collagenous (NC) regions (aa 1-121, 217-269, 442-463, and 700-717) interspersed among three collagenous (COL) domains (aa 122-216, 270-441, and 464-699). Multiple splice forms exist and typically involve deletions of 12-30 aa between aa 220-705. Proteolytic cleavage after Arg108 generates an 85-90 kDa soluble form. Over aa 109-668 (based on a human isoform [NP_542992] that shows a deletion of aa 239-260, 551-565, and 616-627), human COL13A1 shares 91% aa sequence identity with mouse COL13A1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.