

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TOP2B in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human TOP2B aa 1187-1621 Accession # Q02880
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Knockout Validated	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

TOP2B (DNA Topoisomerase II β) is a 180-185 kDa member of the type IIA subfamily, topoisomerase family of molecules. It is ubiquitously expressed, and represents the larger of two known topoisomerases (the 2A/α-form being 170 kDa in size). TOP2A is essential for cell division, while TOP2B is active postmitotically. In an ATP-dependent manner, homodimeric TOP2B relieves torsional stress created in DNA during transcription, or as a consequence of replication. TOP2B first induces cleavage of double-stranded DNA, creating a space that allows for the physical repositioning of chromatin and a reduction in tension. This is followed by closure and ligation of the cleaved ends to recreate the original DNA structure. Human TOP2B is 1626 amino acids (aa) in length. It contains an ATPase domain (aa 101-201), an NES (aa 1034-1044), more than 30 Ser/Thr phosphorylation sites and a C-terminal DTHCT region (aa 1508-1611). There is one splice variant that shows a deletion of aa 24-28. Over aa 1187-1621, human TOP2B shares 91% aa identity with mouse TOP2B.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.