

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CD302/CLEC13A in direct ELISAs and Western blots. In direct ELISAs, approximately 30% cross-reactivity with recombinant human CD302 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse CD302/CLEC13A Asp21-His165 Accession # Q9DCG2
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CD302 (also DCL-1 and CLEC13A) is a 31-36 kDa member of the C-type lectin domain family of proteins. It is expressed on cells involved in phagocytosis, including neutrophils, monocytes, dendritic cells and macrophages. CD302 is suggested to facilitate phagocytosis, and participate in cell adhesion and migration. Mature mouse CD302 is a 208 amino acid (aa) type I transmembrane glycoprotein. It contains a 146 aa extracellular domain (ECD) (aa 21-165) plus a 42 aa cytoplasmic region. The ECD possesses one C-type lectin domain (aa 30-149) that apparently cannot bind carbohydrate. In human, CD302 has also been detected in dendritic cells as the C-terminal portion of a 215 kDa fusion protein that involves DEC205. CD302 contributes aa 23-232 to this fusion protein. Mouse CD302 has two potential splice variants. One shows an alternative start site at Met61, while another shows a deletion of aa 154-162. Over aa 21-165, mouse CD302 shares 82% and 92% aa identity with human and rat CD302, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.