

#### DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Protein O-Glucosyltransferase 1/POGLUT1 in direct ELISAs and Western blots.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human Protein O-Glucosyltransferase 1/POGLUT1 Arg24-Leu388 Accession # Q8NBL1
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below

#### DATA



#### PREPARATION AND STORAGE

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.2 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

#### BACKGROUND

O-Glucosyltransferase1 (POGLUT1) is a homologue of Rumi from Drosophila, an endoplasmic reticulum (ER)-retaining glucosyltransferase, that adds glucose to serine residues within the consensus sequence of C1-X-S-X-P-C2 in Notch EGF repeats, thereby regulating cell-fate decisions (1). It is also known as CAP10-like protein (2) and KTELC1 due to the highly conserved CAP10 domain and the presence of an ER-retaining motif, KTEL, at the C-terminus. The human gene is reported to be involved in the pathogenesis of both acute myelogenous and T-acute lymphoblastic leukemias (3). Recently, POGLUT1 has been demonstrated in a phosphatase-coupled glycosyltransferase assay to have hydrolase activity on UDP-Glc (4).

#### References:

1. Acar, M. *et al.* (2008) *Cell* **132**:247.
2. Teng, Y. *et al.* (2006) *Gene* **371**:7
3. Wang, Y. *et al.* (2010) *Genet Test Mol Biomarkers* **14**:127.
4. Wu, Z.L. *et al.* (2010) *Glycobiology*, in press.