

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human PRKD3/nPKCv in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human PRKD3/nPKCv Ile465-Val559 Accession # O94806
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

PKCv (Protein kinase C- ν ; also PKD3 and PRKD3) is a 105-110 kDa member of the PKD subfamily, CAMK Ser/Thr protein kinase family of enzymes. It is expressed in highly diverse cell types such as pancreatic aciner cells, skeletal muscle cells, B cells and sensory neurons, and it is activated by PKC-mediated phosphorylation. Active PKCv associates with membrane compartments, and appears to participate with VAMP2 in cargo transport and vesicular trafficking. It also acts on class II HDACs in the nucleus, resulting in gene transcription. Human PKCv is 890 amino acids (aa) in length. It contains two DAG-type zinc finger regions (aa 154-204 and 271-321), one PH domain (aa 416-532) and a large protein kinase domain (aa 576-832). There are at least 25 utilized Ser/Thr phosphorylation sites that span the length of the molecule. PKCv possesses one potential splice variant that shows a 17 aa substitution for aa 595-890. Over aa 465-559, human PKCv is 92% aa identical to mouse PKCv.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.