

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects recombinant human IGLC2 in direct ELISAs and Western blots. Detects endogenous Immunoglobulin lambda constant region in natural samples by Western blots.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human IGLC2 Gln2-Ser106 Accession # P0CG05
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 μm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 μg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.2 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Immunoglobulin lambda constant region is a Ke-Oz- component of immunoglobulin (Ig) lambda light chains and is found C-terminal to the joining region of the light chain. IGLC2 is 105 amino acids (aa) in length. It contains one Ig-like domain (aa 7-100) and generates a disulfide bond with an IgH chain via Cys104. One potential variant shows a three aa substitution for aa 7-9. The closest mouse λ ortholog shares 61% aa identity with full-length human IGLC2. IGLC2 shares 93 - 98 % aa identity with IGLC1, IGLC3 and IGLC7.