

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human α -2,8-Sialyltransferase 8B/ST8SIA2 in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human α -2,8-Sialyltransferase 8B/ST8SIA2 Asp24-Thr375 Accession # Q92186
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.
Intracellular Staining by Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

ST8SIA2, also known as sialyltransferase X, is mainly expressed during embryonic development (4) and shows strict preference on NCAM (6). Polysialic acid (PSA), abundant on the neural cell adhesion molecule (NCAM) during embryonic development, acts as an anti-adhesive glycan to negatively modulate the adhesive properties of NCAM (1). PSA expression decreases promptly after birth, and becomes restricted to the hippocampus, hypothalamus, and olfactory bulb, areas of the brain that require continuous cell migration and synaptic plasticity (2). Expression of PSA in cancer cells has been suggested to increase tumor invasiveness and to promote tumor growth (3). The temporal regulation of PSA is dependent on the expression of two polysialyltransferases, ST8SIA4 and ST8SIA2 (4, 5). The high degree of substrate specificity is achieved through specific enzyme-substrate recognition at both the protein sequence and glycan structure levels (6, 7). Like most glycosyltransferases, ST8SIA2 is a Golgi-resident type II membrane protein. The activity of this enzyme has been measured with a phosphatase-coupled method (8).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.