

## DESCRIPTION


<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects human and mouse Peroxiredoxin 3 in direct ELISAs and Western blots. In direct ELISAs, approximately 6% cross-reactivity with recombinant mouse (rm) Peroxiredoxin 2 is observed and less than 1% cross-reactivity with recombinant human (rh) Peroxiredoxin 1, rmPeroxiredoxin 5, and rmPeroxiredoxin 6 is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant mouse Peroxiredoxin 3 Met1-Gln257 Accession # P20108
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Sterile PBS to a final concentration of 0.2 mg/mL.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Peroxiredoxin-3 (Prx-3/III; also AOP-1, MER5 and thioredoxin-dependent peroxidase reductase) is a ubiquitous, 22-28 kDa mitochondrial antioxidant enzyme that belongs to the 2-Cys class of the TSA/ahpC family of peroxiredoxins. Prx-3 is known to act as either a homodimer, or a decamer, and scavenge reactive oxygen species generated by oxidative stress. The mouse Prx-3 precursor molecule is 257 amino acids (aa) in length. It contains a cleavable N-terminal 63 aa mitochondrial targeting sequence, plus a 194 aa mature enzyme that shows a thioredoxin domain between aa 64-222. There are two catalytic cysteines, one at Cys109 and another at Cys230 of the precursor. Prx-3 undergoes a phosphorylation at Thr147 that reduces its activity. One potential splice form shows a deletion of Arg149, Lys150 and Arg185. Full-length mouse Prx-3 is 86% and 95% aa identical to human and rat Prx-3, respectively.