

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse PLA2G4A in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human PLA2G4A Met1-Ala749 Accession # NP_077734
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

PLA2G4A (Phospholipase A2 Group IVA; also cytosolic phospholipase A2 alpha/cPLA2α and lysophospholipase) is an 100-110 kDa member of the cytosolic PLA2 family of enzymes. Expression can be induced in select cell types, including endothelium, smooth muscle, macrophages, PMNs, fibroblasts, mast cells and platelets. Tumor cells often serve as sources for cPLA2α activity. cPLA2α is synthesized in the cytosol where, upon activation, it translocates to sites containing cell membranes, including the Golgi, ER and nuclear membrane. This translocation allows for enzymatic action on membrane phospholipids, generating free arachidonic acid that is converted into proinflammatory eicosanoids. Human cPLA2α is 749 amino acids (aa) in length. It contains a phospholipid-binding and membrane-association C2 domain (aa 1-178), and an overlapping PLA2c domain that demonstrates catalytic activity (aa 140-740). While phosphorylation on Ser505 and Ser727 contribute to activation, there are at least five other utilized phosphorylation sites on the molecule. Phosphorylation may increase the MW of cPLA2α in SDS-Page to about 100 kDa. Full-length (aa 1-749) human PLA2G4A/cPLA2α shares 94% aa sequence identity with mouse PLA2G4A.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.